References

- 1. Kirwan PD, Chau C, Brown AE et al. HIV in the UK. 2016 report. Available at: www.gov.uk/government/publications/hiv-inthe-united-kingdom (accessed May 2017).
- 2. Claymore E. Fact sheet: undetectable viral load. Available at: betablog.org/fact-sheet-undetectable-viral-load (accessed May
- 3. Donnell D, Baeten JM, Kiarie J et al. Heterosexual HIV-1 transmission after initiation of antiretroviral therapy: a prospective cohort analysis. Lancet 2010, 375, 2092-2098.
- 4. Bello KJ, Mesner O, O'Bryan TA et al. Factors associated with 10 years of continuous viral load suppression on HAART. BMC Infect Dis 2016, 16, 351.
- 5. May MT, Gompels M, Delpech V et al. Impact on life expectancy of HIV-1 positive individuals of CD4+ cell count and viral load response to antiretroviral therapy. AIDS 2014, 28, 1193-1202

- 6. Peterson JL, Rintamaki LS, Brashers DE et al. The forms and functions of peer social support for people living with HIV. J Assoc Nurses AIDS Care 2012, 23, 294-305.
- 7. Sayles JN, Ryan GW, Silver JS et al. Experiences of social stigma and implications for healthcare among a diverse population of HIV positive adults. J Urban Health 2007, 84, 814-828.
- 8. Rintamaki LS, Davis TC, Skripkauskas S et al. Social stigma concerns and HIV medication adherence. AIDS Patient Care STDS 2006, 20 359-368
- 9. National AIDS Trust. HIV: public knowledge and attitudes, 2014. Available at: www.nat.org.uk/publication/hivpublicknowledge-and-attitudes-2014 (accessed May 2017).

Correspondence: Irina Lut stigmasurvey@fpa.org.uk

Mental health and HIV: answers to the self-assessment quiz

- 1. The correct answer is **b**
- 2. Answers **a**, **b** and **e** are correct
- 3. Answers **b**, **c** and **e** are correct
- 4. Answers **b** and **d** are correct
- 5. The **sympathetic** nervous system causes release of the 'stress hormone' adrenalin from the **adrenal** glands as a result of perceived threat. This hormone causes an increase in the **heart** and **respiratory rate** as well as gastrointestinal effects such as nausea/
 - vomiting and diarrhoea
- 6. Answer **d** is correct. EDMR stands for **Eye** movement desensitisation and reprocessing.
- 7. Any from the list contained in Box 6 in the article, including: reduced treatment adherence, loss of patient engagement,

- increased use of alcohol and drug use, clinical decline, increased hospital admissions, high-risk sexual behaviour and increased risk of onwards HIV transmission
- 8. Answers **b**, **c** and **d** are correct
- 9. Your answer could include: psychoactive substance use including alcohol, age-related or other neurological disease, e.g. Alzheimer's or Parkinson's diseases, CVD, hypoxia, drug side effects, cytomegalovirus infection, IRIS, hepatitis C, depression and others
- 10. Answers **b** and **d** are correct. Note that all PLWH should be screened for the presence of both mental health problems and cognitive difficulties within 3 months of their diagnosis