

2016



Exploring shared decision making in HIV nursing care

Dr Catrin Evans
University of Nottingham

Ms Michelle Croston
Manchester Metropolitan University

Mr John McLuskey
University of Nottingham

Contents

Executive summary	3
Introduction	4
Study aims and objectives	4
Background	4
Methods	6
Part 1: qualitative study	6
Part 2: online survey	6
Ethics	6
Findings from Part 1: qualitative study	7
Participants	7
Themes	7
Findings from Part 2: online survey	16
Demographic details of the survey participants	16
Interpretation of SDM	17
The role of nurses in SDM	18
Proficiencies for SDM	20
Challenges for SDM	21
Implications for education and training around SDM	22
Discussion	25
Limitations of the research	26
Recommendations	26
Conclusions	26
Acknowledgements	26
Abbreviations	26
References	27
Appendix: online survey	28
Preamble	28
Questions	28
Shared decision making	29

Executive summary

Background

Shared decision making (SDM) is an important part of promoting self-management and empowerment for patients with long-term health conditions. There has been little empirical research on the nature and practice of SDM in HIV care, in spite of national guidelines promoting the approach.

Research aim

This research project aimed to explore current views and practices amongst UK HIV nurses regarding SDM in order to identify training and support needs.

Methods

This was a mixed-methods study. Part 1 was a qualitative study in which four focus-group discussions (15 participants in total) were held to explore HIV nurses' views and practices around SDM. These were thematically analysed and the results were used to develop Part 2, an online survey that was sent to all members of the National HIV Nurses Association (NHIVNA). The survey sought to identify knowledge, challenges, gaps and training needs in relation to SDM. The survey received 64 responses out of a possible 258, a response rate of 25%.

Results

Qualitative data showed that nurses are supportive of SDM and strive to implement it in everyday practice. Nurses understand SDM as a collaborative process but one that must be negotiated not only with the patient but also with the wider MDT. Nurses face several patient-related, organisational and health-system challenges in implementing SDM. The online survey identified a need for more training on SDM (especially in supporting complex patients) and a need for more resources/decision aids to help facilitate SDM.

Conclusions

SDM is an important aspect of nursing care for people living with HIV. Nurses need more training and resources to implement SDM effectively. In order to develop such training and resources and to better understand the meaning of SDM in HIV care, there is a need for research on patient perspectives and experiences in this area.

Introduction

This report presents a mixed-methods research study, exploring the views, practices and experiences of UK HIV nurses in relation to shared decision making (SDM).

The report is presented in two parts, reflecting two different phases of the research. Part 1 presents the results of a qualitative study that used focus-group discussions (FGDs) to explore HIV nurses' views and practices around SDM. The results of this phase were then used to develop Part 2 in which an online survey was sent to all members of the National HIV Nurses Association (NHIVNA). The survey sought to identify knowledge, challenges, gaps and training needs in relation to SDM.

Study aims and objectives

The overall aim of the research was to explore current views and practices amongst UK HIV nurses regarding SDM in order to identify training and support needs.

Specific study objectives were:

- To explore what nurses understand by SDM;
- To explore perceived barriers/facilitators to SDM in HIV care;
- To explore how SDM is practised in everyday HIV clinical settings;
- To explore HIV nurses' views of current SDM resources and to identify what additional resources may be required;
- To identify specific training needs in relation to SDM in HIV care;
- To establish the most acceptable format for additional learning material or skills development in relation to this topic.

Background

People living with HIV are required to make complex and multiple decisions about managing their condition [1]. Shared decision making (SDM) is increasingly advocated as the preferred model for engaging with patients in making choices about their care. A commonly used definition for SDM is provided by Coulter and Collins (p.2) [2].

'Shared decision-making is a process in which clinicians and patients work together to clarify treatment, management or self-management support goals, sharing information about options and preferred outcomes with the aim of reaching mutual agreement on the best course of action ... Shared decision-making explicitly recognises a patient's right to make decisions about their care, ensuring they are fully informed about the options they face. This involves providing them with reliable evidence-based information on the likely benefits and harms of interventions or actions, including any uncertainties and risks, eliciting their preferences and supporting implementation.'

SDM is embraced as a key feature of recent NHS policy whereby patients should be at the heart of all aspects of health-related decision making [3]. Current healthcare policy encourages an increased focus on self-management; indeed, evidence suggests that poor health literacy contributes to suboptimal care, poorer health outcomes for affected individuals and avoidable costs within health systems [3,4]. As such, the philosophy and practice of SDM is concerned with listening to patients' wishes, informing and educating patients about different options, being sensitive to patients' needs, and engaging in an ongoing dialogue to develop mutually agreed appropriate treatment or care pathways [2].

The process of developing SDM between patients and providers is considered particularly necessary to promote patient empowerment for the management of long-term conditions. This approach to care has been well established in other disease areas [5]. A recent review by Stiggelbout *et al.* (p.1173) [6], suggests that there are four clear steps for SDM:

1. The professional informs the patient that a decision is to be made and that the patient's opinion is important;
2. The professional explains the options and the pros and cons of each relevant option;
3. The professional and patient discuss the patient's preferences; the professional supports the patient in deliberation;
4. The professional and patient discuss patient's decisional role preference, make or defer the decision, and discuss possible follow-up.

As HIV moves toward a chronic disease management model, it is important to ensure that SDM principles are also embedded within HIV care [7]. The key areas where people living with HIV face complex decisions have been described in a 2010 systematic literature review as: (i) whether or not to disclose their diagnosis to others; (ii) decisions about adherence to treatments; and (iii) decisions about sexual activity and parenthood [1].

The need for patient empowerment and for patients to be involved in decisions about their healthcare is reflected in the British HIV Association (BHIVA) *Standards of Care for People Living with HIV* [8], particularly Standards 9 and 10. Standard 9 suggests that patients should be involved within their care and self-management as these approaches can ultimately lead to improvements in patients' quality of life. Standard 10 states, '*People living with HIV should be actively involved in decisions relating to their own care and treatment as they wish*'.

The European AIDS Clinical Society (EACS) treatment guidelines also support an SDM approach [9]. They advocate using a 'WEMS' technique (waiting 3 seconds, echoing, mirroring, summarising) to assess readiness to start treatment. This technique is similar to motivational interviewing techniques. This recommendation goes some way to addressing SDM in relation to starting treatment; however, it does not address how patients are involved in decisions throughout the rest of their healthcare journey or for health issues other than treatment adherence.

Within the UK, other than the BHIVA *Standards of Care* [8], there are no suggested guidelines or resources available to help healthcare practitioners facilitate SDM within HIV care. Similarly, whilst considerable research on SDM has been undertaken in other disease areas [5,10–12], very little research has explored SDM specifically in relation to HIV care [13–15].

The goal of this research therefore, was to deliver insights into SDM within HIV care in the UK, from a nursing perspective.

Methods

A two part sequential mixed methods design was adopted [16].

Part 1: qualitative study

The qualitative study component comprised four focus-group discussions with HIV nurses. These took place from February to April 2015. The focus groups were designed to elicit nurses' views and experiences around SDM and to identify areas where SDM may be contested or where it may be particularly challenging [17]. The groups were designed to reflect experiences of nurses in different parts of the country, and were undertaken in four different geographical regions (North West, Midlands, London and the South East).

Recruitment to the focus groups was through regional subgroups within NHIVNA. These are small, regionally focused groups that act as hubs for local communication, training and development around HIV nursing. The Chairperson of each group invited members by email to attend an SDM project focus-group discussion. The inclusion criteria were nurses who were Band 5 or above with HIV patient caseloads or working within HIV services. A total of 15 nurses took part (see Table 1).

Table 1: Focus-group details

Region	Number of participants
North West	2
Midlands	5
London	2
South East	6
Total	15

The focus groups were facilitated by the project research assistant and one member of the project advisory team. Each focus group was audio recorded and transcribed verbatim.

Data were analysed using a thematic analysis approach. Segments of the transcripts were coded according to their meanings [18]. The codes were then grouped and further analysed into several core themes and associated subthemes [19,20]. The project team worked together to develop and refine the emerging interpretations [21].

Part 2: online survey

The themes from the qualitative study were used to inform the design of an online questionnaire that aimed to identify HIV nurses' knowledge, practices and training needs in relation to SDM (see Appendix). The questionnaire was piloted by five nurses in one geographical region and then finalised.

An email with the link to the SDM online survey was sent out to all NHIVNA members via the NHIVNA Secretariat. These were sent in early June 2015 and followed by two reminder emails.

The questionnaire results were analysed using descriptive statistics.

Ethics

Ethical approval for the research was granted by the University of Nottingham, Faculty of Medicine and Health Sciences Research Ethics Committee. Participation in both parts of the study was entirely voluntary. Participation in the focus groups required written consent. Response to the online survey was taken to imply consent. All participant contributions have been anonymised.

Findings from Part 1: qualitative study

Participants

Thirteen of the fifteen nurses who participated in the focus groups filled in a socio-demographic data form, summarised in Table 2.

Table 2: Participant characteristics

Band	Gender	Current job setting	Current job title/ description	Length of time in the area of HIV	Length of time in current role
6	Female	Community	Community Clinical Nurse Specialist	24 years	8 years
6	Female	Community	Clinical Research Nurse	5 years	2 months
7	Male	Community, Specialist HIV Outpatients Department, Genitourinary Medicine, Prison	Clinical Nurse Specialist	28 years	
6	Male	Community	HIV Clinical Nurse Specialist	15 years	7 years
5	Female	Specialist HIV Outpatients Department	Staff Nurse	3 years	
8	Female	Specialist HIV Outpatients Department	Nurse Consultant	15+ years	10 years
7	Male	Community, Specialist HIV Outpatients Department	Senior HIV Nurse Specialist	25 years	10 years
7	Female	Specialist HIV Outpatients Department	Clinical Nurse Specialist women's health, sexual health and HIV, Specialist Practitioner	12 years	2 years
7	Female	Community, Infectious Diseases Unit	HIV Clinical Nurse Specialist	17 years	10 years
7	Female	Community, Specialist HIV Outpatients Department	HIV Clinical Nurse Specialist	23 years	13.5 years
7	Female	Community, Infectious Diseases Unit	HIV Clinical Nurse Specialist	9 years	9 months
7	Female	Community, Genitourinary Medicine, Infectious Diseases Unit	HIV Clinical Nurse Specialist	25 Years	
7	Female	Community	HIV Clinical Nurse Specialist	5 years	5 years

Themes

Four major themes were identified and are outlined in Table 3. These are presented in turn below, with supporting quotations to illustrate key points.

Table 3: Qualitative study themes

Theme	Subthemes
Interpretation of SDM	Varied understandings Varied groups involved Varied focus
The role of the nurse in SDM	Advocate Information giver, teacher, translator Expert Health promoter
Proficiencies for SDM	Knowledge Interpersonal attributes Communication skills
Challenges in SDM	Patient factors Social factors Organisational factors Health system factors

Theme 1: interpretation of shared decision making

A key theme that emerged was the varied understanding that individuals had of shared decision making. There were two key distinctions. The first was SDM primarily as a consultative, **information giving** activity:

'[Shared decision making is] Trying to check out how they feel about what you've said, how they feel about suggestions you've made for care. Whether they're comfortable with that, whether they're satisfied with that. Erm, and just making sure you get their feedback along the way really.' (Nurse 2, Focus Group 1)

The second was a view of SDM more in terms of **partnership** working:

'I suppose I see it as a partnership really, working with the patients and giving them information to help them to make decisions, supporting them in whatever decisions they make. So I see it more as a partnership.' (Nurse 5, Focus Group 2)

All groups, however, agreed that SDM was a **process**. Other differences in perceptions of SDM concerned **who** the decision making process involved. For example, in some cases, SDM was understood as a process of decision making primarily between practitioners and patients, whereas for others, SDM was perceived more as a process of 'shared care', i.e. as a process of discussion amongst a multidisciplinary team (MDT) where decisions, outcomes and plans were then later fed back to the patient. Such MDT discussions were described as typically doctor-led.

'The decision, the consultant and doctors and teams meet, medical teams decide on what needs ... what the patient needs to have.' (Nurse 2, Focus Group 3)

Another aspect of SDM concerned its **focus**. The majority of discussion on SDM referred to treatment options rather than other elements of HIV care. However, some of the groups noted that SDM could be applied to all aspects of care:

'Yes that is what I mean about the more holistic things isn't it? And I mean we're talking a lot about treatment but shared decision making isn't just about treatment ... for some people it is about how often they come to clinic, who it is they're going to see.' (Nurse 4, Focus Group 4)

Taking such a holistic approach to SDM meant that the process was still considered important even when it concerned a patient's refusal to engage in care:

'... when a person actually refuses something it is still part of a choice, it is shared decision making so you cannot force the patient, you have to respect their decision ... So it is good to respect both sides. I think in decision making that the patient has the right to say no and to say yes or to be involved in their care.' (Nurse 3, Focus Group 4)

Theme 2: the nurse's role in SDM

Typically, nurses described complex multi-faceted roles involving interactions with multiple stakeholders across multiple settings. Nurses identified four different functions within their roles that could support SDM.

The first was of 'advocate', speaking for the patient caught in the midst of a complex health system.

'You're an advocate aren't you? As well, even though it's their decision, you're there to advocate for them, so you're listening to their views so you then have to advocate for what they want.' (Nurse 1, Focus Group 1)

Where decisions were being taken in the patient's absence (for example, within an MDT meeting), nurses attempted to facilitate SDM by speaking up for what they thought the patient would want:

'I think we make sure for our patients that we shout up loud don't we? If we're not happy, if we think that the patient wouldn't be happy about that or they have expressed that they're not happy about that, we are prepared to shout up loud on behalf of that patient because they are not there at that meeting and talking about their health.' (Nurse 2, Focus Group 4)

The second role function was that of 'information giver' in order to empower patients to make choices:

'... to give them the knowledge so that they can make the decision of what would happen if you ... if you made different decisions? If you decided to go on this treatment or that treatment or no treatment and so I think that is a big empowering ... that we have got the information because they generally, they might not know. Like what would happen with all the different outcomes.' (Nurse 4, Focus Group 2)

The complexity of information available, however, meant that providing information was not a straightforward activity but also encompassed an explicit teaching role, i.e. it was a nurse's job not just to share information but also to ensure that the patient had understood it:

'If you are wanting patients to make an informed decision, you are not only a nurse you are also a teacher, you are going to teach them about how they are going to have that knowledge to make that shared decision ... so it is very important that you be an informed professional as well as a teacher.' (Nurse 3, Focus Group 4)

The teaching dimension of information giving, however, carried an implicit connotation that there was a 'right' answer to a particular choice. For example, nurses noted that a key part of their job was to achieve optimal clinical outcomes for their patients. Therefore, there was an implicit pressure to help support patients to make choices that would achieve (medically defined) positive outcomes. Nurses noted that when patients made choices that were contrary to the prescribed course of action, there was a temptation to feel that their teaching had not been adequate rather than that this outcome may also be an acceptable consequence of SDM. For example, one nurse stated:

'I personally don't think I will ever feel I have given this person enough information if especially if they are still deciding not to take the medication.' (Nurse 2, Focus Group 3)

Within the information-giving role, some nurses also noted that they acted as a ‘translator’ in order to make the relevant information as accessible as possible. This role was often described as having to translate information that had been provided to patients by other health professionals (particularly doctors) into terms that a patient would understand:

‘Shared means shared to me. ... the person who is having the treatment should have some say in what is actually happening, and often ... they are not empowered or they are not spoken to in a language that is meaningful to them. They haven’t got enough information to make a clear decision sometimes and I think often doctors ... and I am stereotyping doctors as a big group, but they explain in doctor language and think that they’re actually doing that, when the person comes out of the room they say “I don’t know what they are on about you know”.’ (Nurse 2, Focus Group 4)

Conversely, another element of this translation role was in helping doctors to understand patients’ wishes. Nurses felt well placed to do this because of their holistic perspective and the good understanding that they may have acquired over time about a patient’s social and emotional situation:

‘But where that patient had decided to not take treatment, well in the knowledge they will die, and that actually being the end result that they are after. That makes the doctors feel very uncomfortable but again I think that is when the nurses come in, because the doctors are a bit like oh they don’t want care, but no that is not what they’re saying. They don’t want to continue on treatment and not wanting to continue on treatment and not wanting care are two very different things aren’t they? They still did want to be looked after and monitored but they did not want to take tablets and they wanted to die.’ (Nurse 5, Focus Group 4)

A third role function identified participants as ‘experts’, having access to specialist knowledge and expertise that gave them a privileged status in suggesting options for patients.

‘Being able to engage patients is the heart of the process of decision making, and number one is being an expert in the field ... to ensure that the patient has confidence ... to give them all the information required to make this, you know decision.’ (Nurse 1, Focus Group 3)

Some nurses drew comparisons between their work and that of other experts, for example a car mechanic. Implicit in this representation was the notion that, as experts, they were best placed to advise an optimal course of action for the patient:

‘... in shared decision making because you know it is supposed to help the patient understand what there is to understand – but at the end of the day ... if I go to see a mechanic about my car, and they say it needs a new pump and I won’t say well I will just have a look myself and check that out will I? I will say OK so what am I ... you know what choices have I got? I haven’t got a choice of ... if I want to run this car I need that new pump so I could have the special one at £200 or I could have a slightly cheaper version, what do you think is the difference between these two? And I will just go yes or no won’t I?’ (Nurse 1, Focus Group 4)

Participants identified a fourth aspect of their role in SDM as ‘health promoter’. Within this role, they described a professional and moral duty to follow clinical guidelines and to promote decisions that would yield (medically and economically defined) good clinical outcomes. As noted above, an SDM ethos sometimes conflicted with this health-promoter role, especially when patients acted in ways that were contrary to clinical guidelines or clinical advice:

‘So our role as well as looking after the general care of the patient is to encourage them to be able to take that medication which we know works ... Then we do have the pressures in the NHS now, I mean if you look at the Standards of Care for People Living

with HIV as well as long-term chronic diseases, we have to avoid hospital admissions – so you find that in a way, this is at the back of your mind. You know what is best for the person and you really would like them to avoid being an inpatient and interrupting a lot of other things in their life.’ (Nurse 2, Focus Group 3)

Participants noted that all four role functions were crucially dependent on being able to develop trusting relationships with their patients, so that patients would recognise their expertise, follow their advice and feel able to express their views and concerns:

‘What helps is when you trust that person and it is like ... if they see a doctor, but then they come away and then talk to us and we have a different relationship with the patient so we can like validate that this doctor’s opinion is a good opinion, it is then ... I think people feel like they are making more of an informed decision about it. That trusting relationship ... you know, you can’t put a price on that because, you know, ‘cause they’ll actually open up and share things that they may not with someone else.’ (Nurse 1, Focus Group 4)

In sum, the role of the nurse in SDM was highly varied. In some cases, nurses sought to promote SDM by acting as advocate for patients. In other situations, the nurses’ role in SDM was more ambiguous, particularly where nurses (as information givers, experts and promoters of health) felt that the optimal course of action was clear.

Theme 3: proficiencies for SDM

Generally, nurses felt that SDM was something that could not be taught but that developed with experience over time. A range of attributes and skills was identified that could facilitate SDM (described below); however, participants noted that knowledge was a key underpinning requirement for these to be put into practice. Knowledge was seen as essential in terms of providing accurate information to patients, but also in terms of being able to present care and treatment options in an individualised way:

‘I think you’ve got to start from the point of view of knowledge, so, you have to understand, every decision that that patient makes, what’s the impact on their care? And you’ve got to be able to explain to them what that impact is, negative or positive ... otherwise, they can’t make informed choices. So you’ve got to have that strong confident knowledge.’ (Nurse 2, Focus Group 1)

In terms of attributes, nurses noted that SDM required them to be flexible, perceptive, non-judgemental and understanding. These attributes were collectively referred to as ‘intuition’ – an ability to read between the lines, to assess non-verbal behaviour and to discern a clearer understanding of what a patient may be feeling:

‘I think using your senses, all of your senses is quite important because sometimes what is coming out of a patient’s mouth isn’t often what they are feeling or they are thinking and by looking at them and listening to them, and sometimes touching them you can tell that actually that isn’t the route that they want to go down, that isn’t really what they’re feeling is their decision and I think using all of those skills eventually leads to quite good intuition and I think that often when professionals follow their intuition it is often the right thing for the patient.’ (Nurse 4, Focus Group 4)

Nurses’ holistic perspective was also mentioned as essential in facilitating SDM:

‘I think from a nursing point of view it is looking at the whole person. Looking at all elements as well. That is a big skill actually isn’t it?’ (Nurse 2, Focus Group 2)

Within all the focus groups, participants were unanimous that the key skill required for SDM was communication. In particular, participants identified listening as an essential skill for delivering

patient-centred care. In addition, it was important to check that patients had understood the nurse by reflecting back on the patient's feelings.

'So definitely – extended communication. Understanding the patient, understanding their ability to comprehend so the nurse has to be skilled in listening, skilled in reading, knowledgeable and also know when they're not making headway. It's the usual, the open questions, the you know, the reflecting back on the patients feelings, the listening skills ... trying to understand um, the historical perspective for some patients as to their attitudes towards tablets in the first place. What it means to them. So it is the more advanced counselling skills.' (Nurse 1, Focus Group 1)

Motivational interviewing was raised in each of the focus groups as an important communication technique. Nurses highlighted its perceived potential to improve the delivery of communication and care in HIV treatment:

'I think also, again I don't know about the rest of you, but maybe using a bit of motivational interviewing, just to see where patients are situated with that particular decision at the time. I've found that useful.' (Nurse 5, Focus Group 2)

However, although many participants felt that motivational interviewing was an essential skill to have, several noted that they themselves did not feel confident in using it:

'I would love to have training in motivational interviewing. I think it would be beneficial for all of our team to have it, I know some people have but, you know, it is only just bits I have picked up, whatever you know, but I think it would be good for nurses to get trained in that.' (Nurse 4, Focus Group 2)

In sum, participants noted that a combination of nursing knowledge, interpersonal attributes and communication skills was required to practise SDM.

Challenges in SDM

There was much discussion in the groups about the difficulties of trying to accomplish SDM in practice. Four main challenges were identified by the participants: patient factors, social factors, organisational factors and health system factors.

In terms of patient factors, participants noted that, with some patient groups, language barriers posed challenges for SDM, both in terms of lengthening the time required within a consultation and also the challenge of working with translators:

'Understanding yeah, again, it's when you use interpreters and when you don't, um, interpreters are not ideal, um, the emotional stuff is almost impossible sometimes, um and I had one lady, she, doesn't think there's a communication problem. She thinks that I understand her perfectly and she understands me so why do I need an interpreter? When I know that she's missing a whole lot and I know that I'm struggling to completely understand everything that she's saying.' (Nurse 2, Focus Group 1)

Participants talked at length about patients who were 'complex' or who had complex needs. These were often related to drug and alcohol problems, mental health problems, social problems (e.g. homelessness or poverty) or medical problems (e.g. multiple co-morbidities). Nurses noted that SDM could be more difficult with such 'complex' patients. For example, in some cases, assessing a patient's capacity to make decisions could be difficult, especially in situations where a patient had opted out of treatment, ceased treatment or chosen not to engage in care:

'If someone is an intravenous drug user or ex-intravenous drug user, already has nowhere to live, chaotic lifestyle, there are problems associated with that and different kind of resources needed to associate and support that person in their decision making and also we don't even know the capacity at the time when they are speaking with you, their

capacity whether they understand, they appreciate what is going on and they want to cooperate with you.' (Nurse 2, Focus Group 3)

In situations where patients had mental health or social problems, HIV was sometimes not a priority for an individual, leading to poor adherence, risky behaviour or missed appointments:

'It is my experience as well, because partly the demographics I see more of that kind of patient group who really struggle with many other issues in their lives that HIV and erm ceases to be a priority and they feel it is just like an encumbrance, kind of, and then, I don't know, maybe partly as well because sometimes it is, people are coming from difficult circumstances.' (Nurse 2, Focus Group 3)

Although a number of nurses expressed their frustration in trying to engage with, and support, such patients, they noted that they none the less tried to adopt a patient-centred approach to care:

'You take every individual that walks through the door, they walk through with whatever they walk through with. Hang ups, no hang ups ... issues, no issues, you just, you just go with wherever they're at don't you? Yeah, absolutely ... and it doesn't matter whether its alcohol that's their problem or whether its em, life style or whether it's just their childhood or their personality. It doesn't matter what it is that might be a barrier to good healthcare you just deal with whatever comes up.' (Nurse 2, Focus Group 1)

Other patient-related factors included religious views and beliefs that suggested paths to healing that were contrary to medical advice. In these situations, nurses discussed how they tried to accept and respect patients' choices:

'I mean you see it a lot with people who have got very strong religious beliefs don't you? You know they can ... it can be very challenging to help look after them in a way that is conducive to good health but that is part of their decision isn't it? And that you know if they think that God is going to kill them or heal them, then we have to work with them and not against that and we have to follow a bit of a journey until they reach a place that they feel either they are going to carry on with that and not take medicines or they are going to take medicines.' (Nurse 4, Focus Group 4)

However, nurses also described feeling frustrated or uncomfortable with some of the choices and behaviour of patients, especially when patients were seen to be taking risks or making poor or uninformed choices with regard to their care:

'... the other thing is the essence of why we went in to the profession which we went in to, why we went in to nursing? You were challenged by that patient who says I choose not to get well, I choose to die, I choose not to take the medication. For me that is where the discomfort I think emanates from ... But then how do we allow, how do we get over our own feelings of being like to borrow the word paternalistic and say you have to do this, you have ... it is a difficult kind of area because we went in to nursing to make people better, to try and help them whether it is by personal care, doing the injections ... and then now you're being challenged by the person's decision to say actually I am not going to have this.' (Nurse 2, Focus Group 3)

The topic of 'avoidable death' was something that came up in each of the focus group discussions as one of the most challenging issues to address within an SDM approach. The nurses reflected at length about the evolution of HIV treatment and its transformation into a manageable long-term condition. Thereby, when patients decided not to engage in treatment or HIV care, the nurses saw this as unnecessarily making the decision to die:

'They stop taking their tablets and then that can be quite a challenge for health professionals who are thinking you're going to die without these and yet ultimately the person feels they can't do it and we're thinking look it is simple, just put them in your

mouth and take them but it isn't that simple is it? I think the challenge for us as healthcare providers is saying if someone's internal choices are different to ours, we can't override that and therefore the shared collaboration is then you know what? I have to go with you on your journey, if you're going to die I will still offer the best care that we have available.' (Nurse 4, Focus Group 4)

In terms of social factors, all groups raised the issue and impact of stigma on SDM. Nurses described stigma as something that was still very much a part of HIV and was seen as a barrier to SDM, primarily because fear and stigma prevented patients from accepting or disclosing their diagnosis and engaging with treatment:

'Sadly, the stigma is still very much there and the issues around disclosure and sharing it with anybody is still a massive issue.' (Nurse 2, Focus Group 1)

Stigma was described not only as a problem in terms of patients' personal lives or reluctance to engage, but was also identified as a challenge in terms of the treatment patients received from other healthcare professions. Nurses noted that patients were often referred back to the HIV department for the main part of their care, even if other departments should have been taking the lead. This was attributed to ongoing stigma from other nurses and a lack of confidence amongst other professionals in managing HIV. Together, they posed challenges in terms of achieving an SDM approach within a shared care model:

'We offer some training to practice nurses and stuff and the ignorance that is still around is unbelievable. The nurses, just, I mean, I know from colleagues from like when I was district nursing, how their opinion of HIV, and these are professionals ... So, with shared care, I think we are always going to have to work towards it and I think the stigma will be a hindrance. I don't personally think the stigma will go in our lifetime but just to keep chipping away at it and just keep moving forward on it.' (Nurse 2, Focus group 2)

In terms of organisational factors, a key challenge was trying to engage with SDM under time constraints, for example, short consultation times, or managing very complex patients who required more time than usual. SDM was perceived as lengthening the time required for consultations, and that without providing additional time, SDM was difficult to implement in practice:

'Time, you've gotta have time, you've gotta give time to the patient. The more shared it is I think the longer the time that is needed to go through that process so I think time is a big factor in that you know appointments are a ticking clock aren't they? You know you have got to get them through the clinic so I think time ... our clinic at times you know I wouldn't envy them sitting there that length of time so both from the professionals point of view but from the client, patients point of view time is a big factor.' (Nurse 2, Focus Group 4)

Organisational factors also placed constraints on developing services that could be more flexible or accessible and would therefore enable better patient engagement and SDM. For example, it was suggested that the ability to work in the community or to provide late or early appointments, or to follow up non-attendees was important but not always possible.

'With us in community, that's what we tend to do, the nurses in the hospital might phone us and say such and such has DNA'd their appointment ... and then you know, if they're known to us, we might try and get hold of them or we'll just go round ... on the off chance if you know it's safe ... In a clinic setting because of course when I was based in a clinic setting we didn't have those kind of opportunities and that is why I feel such a privilege to be in the community setting at the moment.' (Nurse 2, Focus Group 2)

Participants also identified several more macro-level health-system-related challenges to SDM. One of the major factors was that HIV care, like other areas of care, was being delivered in a context of local, national and international targets, standards and guidelines. All departments were under pressure to achieve targets around testing, treatment initiation, treatment adherence and viral loads.

Participants noted that whilst targets could be positive, they also created an implicit pressure 'within the system' to guide patients towards particular HIV care choices, which could create tensions for an SDM approach:

'So if you have got global goals and you are like 90/90/90 so it tells the whole world are going to get everyone tested, everyone on treatment, every undetectable and then HIV doesn't spread anymore, it is the miracle fulfilled. And then we're working with individuals in you know very personal ways and they are telling us very personal things ... but we're also trying to meet those targets. All I am motivated to do is get you on treatment or in a study or to come at every agreed interval. I think that is quite a challenge for nurses ... Public Health England and all the people who we report to, we are judged against someone who fails and even by how many people in your clinic are undetectable – you know – you feel in the middle.' (Nurse 2, Focus Group 3)

Similarly, nurses noted that budgetary considerations placed constraints on patient choice of treatment options. Hence, if patients wanted to switch their medication or choose not to opt for certain medication, this could be a problem for the nurses in delivering SDM:

'These are the choices, but the choices are constrained by the London Consortium or whatever deciding about you know, yes a patient can choose whether they go on or off treatment if they want to but there are some limitations ... Also, the fact that we're using maybe slightly different drugs you know that someone saying I want that one pill, well it is actually well, we're not prescribing that one pill so much now we are prescribing it in two because it is cheaper.' (Nurse 1, Focus Group 3)

In conclusion, SDM was something that nurses felt they did on a day-to-day basis. None the less, the issues raised within this section show that nurses face multiple interlinked challenges in enacting SDM in practice.

Findings from Part 2: online survey

This section presents the survey participants' demographic details followed by the main survey findings. These are categorised into five thematic areas, following the same themes arising from the qualitative study, but including one additional theme about suggestions for education and training.

Each survey question is presented below in the format in which it was asked in the questionnaire. The results for each question are summarised in a table followed by some short explanatory text.

The online survey was sent by email to NHIVNA's membership (258 members). Sixty-four responses were received, constituting a response rate of 25%.

Demographic details of the survey participants

Q.1 Nurses' place of work

Work setting	Response (n = 64)
Specialist HIV outpatients department	37.5% (n = 24)
Genitourinary medicine	20.3% (n = 13)
Other (e.g. non-clinical/advisory/research and managerial roles)	17.2% (n = 11)
Community	10.9% (n = 7)
Infectious diseases unit	7.8% (n = 5)
Specialist HIV inpatient care	4.7% (n = 3)
Midwifery/obstetrics	1.6% (n = 1)

The majority of respondents (37.5%) worked in specialist HIV outpatient departments. Thirteen nurses (20.3%) worked in genitourinary medicine and seven nurses (10.9%) worked in the community.

Q.2 Geographical region

Geographical area	Response (n = 64)
England	85.9% (n = 55)
Scotland	9.4% (n = 6)
Wales	4.7% (n = 3)
Northern Ireland	0% (n = 0)

The majority of the respondents (85.9%) worked in England.

Q.3 Size of patient cohort

Size of cohort	Response (n = 64)
Up to and including 500	53.1% (n = 34)
Over 501	46.9% (n = 30)

The approximate size of the HIV patient cohort in each individual’s work setting was almost even, with 34 nurses having a cohort of 500 or fewer patients and 30 with a cohort of 501 or more patients.

Q.4 Current band of employment

Band	Response (n = 64)
Band 8	14.1% (n = 9)
Band 7	46.9% (n = 30)
Band 6	31.3% (n = 20)
Band 5	7.8% (n = 5)

All nurses who took part in the survey were NHS employed and fairly experienced. Most nurses were Band 7 (46.9%) and Band 6 (31.3%).

Q.5 Length of time working in HIV care

Time	Response (n = 64)
More than 10 years	62.5% (n = 40)
5–10 years	18.8% (n = 12)
2–5 years	7.8% (n = 5)
1–2 years	4.7% (n = 3)
Less than 1 year	4.7% (n = 3)
Not specialising in this field	1.6% (n = 1)

The majority of nurses (62.5%) who took part in the survey had worked specifically in HIV care for over 10 years.

Interpretation of SDM

Q.6 In your view, which of the following statements best defines shared decision making in HIV nursing care? (Please select one.)

Definition	Response (n = 64)
A collaborative process that allows patients and their providers to make healthcare decisions together	92.2% (n = 59)
Giving the patient all the necessary information so that they can make the decision on their own	3.1% (n = 2)
Multidisciplinary teams working together to make the best healthcare decisions for the patient	3.1% (n = 2)
Checking what the patient understands and how the patient feels about what you’ve said	1.6% (n = 1)

Definition	Response (n = 64)
Being an advocate for patients when you know what they want	0% (n = 0)
Planned care which the patient consents to	0% (n = 0)
Shared care management with other health professionals	0% (n = 0)
Informing the patient of the best treatment options available	0% (n = 0)
A process to help engage with a patient when they have disengaged or stopped treatment	0% (n = 0)
Shared decision making is not possible because patients are always asking me what I would do	0% (n = 0)

When asked to select a statement that best defined shared decision making in HIV nursing care, the majority by far of respondents (92.2%) selected ‘A collaborative process that allows patients and their healthcare providers to make healthcare decisions together’.

The role of nurses in SDM

Q.7 In your day-to-day practice, in which activities do you most commonly undertake shared decision making with your client? Please give your answer on the scale of 1 to 5, with 1 indicating ‘very rarely’ and 5 indicating ‘very common’.

Activities	1	2	3	4	5
Adhering to treatment	1.6% (n = 1)	0% (n = 0)	7.8% (n = 5)	23.4% (n = 15)	67.2% (n = 43)
HIV testing	28.1% (n = 18)	12.5% (n = 8)	4.7% (n = 3)	9.4% (n = 6)	45.3% (n = 29)
Sexual risk taking	3.1% (n = 2)	1.6% (n = 1)	23.4% (n = 15)	31.3% (n = 20)	40.6% (n = 26)
Starting HIV treatment	15.6% (n = 10)	7.8% (n = 5)	20.3% (n = 13)	21.9% (n = 14)	34.4% (n = 22)
Switching HIV treatment	20.3% (n = 13)	15.6% (n = 10)	18.8% (n = 12)	15.6% (n = 10)	29.7% (n = 19)
Stopping HIV treatment	34.4% (n = 22)	17.2% (n = 11)	20.3% (n = 13)	20.3% (n = 13)	7.8% (n = 5)
Opting out of treatment	29.7% (n = 19)	15.6% (n = 10)	29.7% (n = 19)	18.8% (n = 12)	6.3% (n = 4)
Psychological care	3.1% (n = 2)	3.1% (n = 2)	15.6% (n = 10)	31.3% (n = 20)	46.9% (n = 30)
Smoking cessation	4.7% (n = 3)	10.9% (n = 7)	32.8% (n = 21)	31.3% (n = 20)	20.3% (n = 13)
Family/pregnancy planning	20.3% (n = 13)	17.2% (n = 11)	23.4% (n = 15)	26.6% (n = 17)	12.5% (n = 8)
Recreational drug use	3.1% (n = 2)	7.8% (n = 5)	28.1% (n = 18)	21.9% (n = 14)	39.1% (n = 25)
Managing HIV treatment along with treatment for other medical conditions	4.7% (n = 3)	14.1% (n = 9)	21.9% (n = 14)	21.9% (n = 14)	37.5% (n = 24)

Activities	1	2	3	4	5
Disclosure of status to others	6.3% (n = 4)	9.4% (n = 6)	31.3% (n = 20)	32.8% (n = 21)	20.3% (n = 13)
Dietary advice	1.6% (n = 1)	15.6% (n = 10)	21.9% (n = 14)	32.8% (n = 21)	28.1% (n = 18)
Exercise advice	1.6% (n = 1)	15.6% (n = 10)	23.4% (n = 15)	32.8% (n = 21)	26.6% (n = 17)
Cardiovascular risk advice	4.7% (n = 3)	12.5% (n = 8)	25% (n = 16)	26.6% (n = 17)	31.3% (n = 20)
Alcohol consumption advice	3.1% (n = 2)	10.9% (n = 7)	14.1% (n = 9)	37.5% (n = 24)	34.4% (n = 22)

Nurses were asked in which activities shared decision making was commonly undertaken. The most common areas were around treatment adherence, HIV testing, psychological care and sexual risk taking. Again, the areas where SDM was least likely to be reported were in stopping, switching or opting out of treatment.

Q.8 What strategies do you most use to facilitate shared decision making? Please rank where 1 = never use and 5 = always use.

Strategies	1	2	3	4	5
General discussion	0% (n = 0)	1.6% (n = 1)	6.3% (n = 4)	43.8% (n = 28)	48.4% (n = 31)
Motivational interviewing	15.6% (n = 10)	15.6% (n = 10)	31.3% (n = 20)	31.3% (n = 20)	6.3% (n = 4)
Use of written decision aids	25% (n = 16)	10.9% (n = 7)	29.7% (n = 19)	21.9% (n = 14)	12.5% (n = 8)
Use of web-based resources	7.8% (n = 5)	12.5% (n = 8)	25% (n = 16)	45.3% (n = 29)	9.4% (n = 6)
Drawing on your own professional knowledge/experience	0% (n = 0)	1.6% (n = 1)	12.5% (n = 8)	43.8% (n = 28)	42.2% (n = 27)
Advocacy with other services or professionals	0% (n = 0)	3.1% (n = 2)	7.8% (n = 5)	62.5% (n = 40)	26.6% (n = 17)
Signposting to other services	0% (n = 0)	1.6% (n = 1)	15.6% (n = 10)	56.3% (n = 36)	26.6% (n = 17)
Linking to peer support	6.3% (n = 4)	4.7% (n = 3)	21.9% (n = 14)	40.6% (n = 26)	26.6% (n = 17)

Nurses were asked about eight different strategies that they might use to facilitate shared decision making. The most commonly used strategies were: signposting to other services, general discussion, drawing on their own professional knowledge and advocacy with other services or professionals.

Q.9 What would help you to implement shared decision making in your day-to-day care? Please tick your top three.

Facilitation of shared decision making	Response
Treatment decision aids	51.6% (n = 33)
More time with patients in consultations	51.6% (n = 33)
More support from other professionals	48.4% (n = 31)
Better background knowledge	46.9% (n = 30)
More educational resources for patients	45.3% (n = 29)
More training in communication skills	28.1% (n = 18)
More support from the voluntary sector	15.6% (n = 10)
None of the above	3.1% (n = 2)
Other	9.4% (n = 6)

Nurses were asked to select three options that would help them to implement shared decision making in their day-to-day care delivery. The top three choices were treatment decision aids (51.6%), more time with patients in consultations (51.6%) and more support from other professionals (48.4%). Several respondents (45.3%) noted that a greater range of educational resources for patients would be helpful. Almost half of the sample (46.9%) noted that better background knowledge would be useful, and 18 respondents (28.1%) felt that more training would help.

Proficiencies for SDM

Q.10 On a scale of 1 to 5, please indicate how confident you feel in undertaking shared decision making in the following areas of care, where 1 = not confident at all and 5 = very confident.

Activities	1	2	3	4	5
HIV testing	4.7% (n = 3)	1.6% (n = 1)	4.7% (n = 3)	14.1% (n = 9)	75% (n = 48)
Adhering to treatment	1.6% (n = 1)	0% (n = 0)	4.7% (n = 3)	31.3% (n = 20)	62.5% (n = 40)
Starting HIV treatment	3.1% (n = 2)	6.3% (n = 4)	17.2% (n = 11)	25% (n = 16)	48.4% (n = 31)
Switching HIV treatment	6.3% (n = 4)	15.6% (n = 10)	25% (n = 16)	25% (n = 16)	28.1% (n = 18)
Stopping HIV treatment	7.8% (n = 5)	14.1% (n = 9)	31.3% (n = 20)	25% (n = 16)	21.9% (n = 14)
Opting out of treatment	9.4% (n = 6)	7.8% (n = 5)	31.3% (n = 20)	31.3% (n = 20)	20.3% (n = 13)
Smoking cessation	0% (n = 0)	6.3% (n = 4)	23.4% (n = 15)	42.2% (n = 27)	28.1% (n = 18)
Sexual risk taking	1.6% (n = 1)	3.1% (n = 2)	17.2% (n = 11)	28.1% (n = 18)	50% (n = 32)
Family/pregnancy planning	6.3% (n = 4)	17.2% (n = 11)	28.1% (n = 18)	32.8% (n = 21)	15.6% (n = 10)
Recreational drug use	3.1% (n = 2)	9.4% (n = 6)	31.3% (n = 20)	39.1% (n = 25)	17.2% (n = 11)
Managing HIV treatment along with treatment for other medical conditions	3.1% (n = 2)	9.4% (n = 6)	31.3% (n = 20)	35.9% (n = 23)	20.3% (n = 13)

Activities	1	2	3	4	5
Psychological care	1.6% (n = 1)	4.7% (n = 3)	18.8% (n = 12)	42.2% (n = 27)	32.8% (n = 21)
Disclosure of status to others	3.1% (n = 2)	3.1% (n = 2)	17.2% (n = 11)	48.4% (n = 31)	28.1% (n = 18)
Dietary advice	1.6% (n = 1)	1.6% (n = 1)	29.7% (n = 19)	39.1% (n = 25)	28.1% (n = 18)
Exercise advice	1.6% (n = 1)	1.6% (n = 1)	23.4% (n = 15)	45.3% (n = 29)	28.1% (n = 18)
Cardiovascular risk advice	1.6% (n = 1)	3.1% (n = 2)	20.3% (n = 13)	43.8% (n = 28)	31.3% (n = 20)
Alcohol consumption advice	0% (n = 0)	4.7% (n = 3)	10.9% (n = 7)	50% (n = 32)	34.4% (n = 22)

Nurses were asked to scale how confident they felt in undertaking shared decision making in 17 different areas. Respondents were most confident in the areas of HIV testing and treatment adherence, but expressed high levels of confidence in most areas listed. Respondents were least confident in the areas of treatment cessation or treatment switching.

Challenges for SDM

Q.11 How easy is it to practise shared decision making in your area of HIV care?

Level	Response (n = 64)
Easy	46.9% (n = 30)
Moderate	42.2% (n = 27)
Very easy	7.8% (n = 5)
Very hard	3.1% (n = 2)
Hard	0% (n = 0)

When asked how easy it was to practise shared decision making in their area of HIV, 30 nurses (46.9%), thought that it was easy and 27 nurses (42.2%) thought that it was moderate.

Q.12 The following have been identified as potential barriers to implementing shared decision making in everyday consultations. Please tick the top three that have applied to you in your clinical role.

Potential barriers	Response
Not enough consultation time	45.3% (n = 29)
Cultural issues/beliefs of patients	45.3% (n = 29)
Patient doesn't want to input into the process/patient wants me to make the decision	39.1% (n = 25)
Lack of up-to-date background knowledge on the patient	21.9% (n = 14)
Lack confidence in my own skills	21.9% (n = 14)
Feeling unskilled	20.3% (n = 13)

Potential barriers	Response
Needing interpreters	17.1% (n = 11)
Not knowing patient adequately	15.6% (n = 10)
Lack of support from colleagues	14% (n = 9)
Treatment guidelines/options available	14% (n = 9)
Organisational targets	14% (n = 9)
The setting doesn't support shared decision making	9.4% (n = 6)
Public Health England targets	7.9% (n = 5)
This is the role of another member of the MDT	7.9% (n = 5)
Other	6.2% (n = 4)

A selection of potential barriers to shared decision making were listed. Nurses were asked to select the top three barriers that had applied to their clinical roles. Two barriers were jointly ranked as most significant: 'not enough consultation time' (45.3%) and 'cultural issues/patient beliefs' (45.3%). The next most common barrier was 'patient doesn't want to input into the process/patient wants me to make the decision' with 39.1% choosing this option.

Q.13 *Are there any limitations of shared decision making? Please select all that apply.*

Limitations	Response
Not everyone wants shared decision making	64% (n = 41)
Not everyone is good at shared decision making	32.8% (n = 21)
There are no limitations to shared decision making	18.8% (n = 12)
Shared decision making takes too much time	10.9% (n = 7)
Shared decision making is not compatible with clinical practice guidelines	6.2% (n = 4)
Other	6.2% (n = 4)

When asked about potential limitations of shared decision making, respondents were given five options to choose from. Forty-one nurses (64%) chose 'not everyone wants shared decision making' and 21 nurses (32.8%) said 'not everyone is good at shared decision making'.

Implications for education and training around SDM

Q.14 *Have you had any specific training on the topic of 'shared decision making'?*

Had any training?	Response (n = 64)
No	85.9% (n = 55)
Yes	9.4% (n = 6)
Unsure	4.7% (n = 3)

Most nurses (85.9%) reported that they had not had any specific training on the topic of shared decision making. Six nurses said that they had had specific training around SDM. This was reported to be, variously, part of a degree course, part of motivational interviewing training or part of a non-medical prescribing course.

Q.15 Please select the top three issues from the options below where greater knowledge/skills would help to enhance your ability to practise shared decision making in HIV care.

Top three issues	Response
Chaotic patients	57.8% (n = 37)
Patient non-attenders	51.6% (n = 33)
Patients with mental health problems	46.9% (n = 30)
Patients who refuse treatment	39% (n = 25)
Patients who take recreational drugs	32.8% (n = 21)
Patients with drug/alcohol addictions	28.7% (n = 18)
Resources that address language barrier	21.9% (n = 14)
Assessing patients' capacity to make treatment decisions	21.9% (n = 14)

Nurses were asked to select their top three issues from eight options, to show the situations in which they felt that greater knowledge/skills would help to enhance their ability to practise shared decision making. The top three responses all referred to patients with complex needs: 'chaotic patients' (57.8%), 'patient non-attenders' (51.6%) and 'patients with mental health problems' (46.9%).

Q.16 Which resources have you accessed to date, in order to enhance your knowledge of shared decision making? Please tick any that apply.

Resources accessed to date	Response
Study days	54.7% (n = 35)
Conference lectures	51.6% (n = 33)
Online resources	43.8% (n = 28)
Medical/nursing journal articles	40.6% (n = 26)
In-house education	39% (n = 25)
Information booklets	28.1% (n = 18)
None	15.6% (n = 10)
Other: please describe	1.6% (n = 1)

Seven types of educational resource options were listed. Nurses were asked which they had accessed to date in order to improve their knowledge of shared decision making. The three most accessed resources were study days (54.7%), conference lectures (51.6%) and online resources (43.8%).

Q.17 In your view, what knowledge/skills would enable you to undertake shared decision making more effectively with your clients? Please select the top three.

What knowledge/skills would enable you to undertake shared decision making more effectively	Response
Greater knowledge of SDM process	78.1% (n = 50)
Access to resources to aid SDM	71.9% (n = 46)
Practical training in SDM process including role playing	56.2% (n = 36)
Motivational interviewing	42.2% (n = 27)
Counselling skills	20.3% (n = 13)
Other	3.1% (n = 2)

Nurses were asked to select three facilitating factors that would enable them to undertake shared decision making more effectively with their patients. The top three chosen were 'greater knowledge of SDM process' (78.1%), 'access to resources to aid SDM' (71.9%) and 'practical training in SDM process including role playing' (56.2%).

Q.18 If further training and education were made available on the subject of shared decision making, how would you prefer to access this? Please give your order of preference, with 1 being your first choice.

Rank further training/education	1	2	3	4	5	6	7	8
e-learning	39.7% (n = 23)	12.1% (n = 7)	10.3% (n = 6)	6.9% (n = 4)	12.1% (n = 7)	3.4% (n = 2)	3.4% (n = 2)	12.1% (n = 7)
Reading materials online	10.7% (n = 6)	23.3% (n = 13)	17.9% (n = 10)	7.1% (n = 4)	10.7% (n = 6)	10.7% (n = 6)	17.9% (n = 10)	1.6% (n = 1)
Hard copy reading materials	14.8% (n = 8)	7.4% (n = 4)	18.5% (n = 10)	13% (n = 7)	14.8% (n = 8)	18.5% (n = 10)	9.3% (n = 5)	3.7% (n = 2)
Study days	35% (n = 21)	20% (n = 12)	13.3% (n = 8)	13.3% (n = 8)	5% (n = 3)	5% (n = 3)	5% (n = 3)	3.3% (n = 2)
Conference lectures	14.3% (n = 8)	17.9% (n = 10)	10.7% (n = 6)	14.3% (n = 8)	16.1% (n = 9)	8.9% (n = 5)	10.7% (n = 6)	7.1% (n = 4)
Interactive workshops at conferences	25.5% (n = 12)	14.5% (n = 8)	18.2% (n = 10)	7.3% (n = 4)	3.6% (n = 2)	10.9% (n = 6)	10.9% (n = 6)	9.1% (n = 5)
Short education sessions run in my workplace	26.3% (n = 15)	21.1% (n = 12)	15.8% (n = 9)	8.8% (n = 5)	5.3% (n = 3)	3.5% (n = 2)	15.8% (n = 9)	3.5% (n = 2)
Other: please describe	12.5% (n = 1)	0% (n = 0)	12.5% (n = 1)	12.5% (n = 1)	0% (n = 0)	12.5% (n = 1)	12.5% (n = 1)	50% (n = 4)

In terms of training and education, nurses were asked their preferences for accessing further training and education on SDM. Respondents were given eight choices and asked to rank each option with 1 being the first choice and 8 being the last choice. Most respondents chose e-learning as their first choice (39.7%). Twenty-one nurses (35%) chose study days as their first choice and 15 (26.3%) selected 'short educational session run in-house'.

Discussion

The two parts of this research study show that HIV nurses are highly supportive of SDM and strive to implement it in their day-to-day activities. For example, in the questionnaire (Q.6), 92% agreed with the statement that SDM is ‘a collaborative process that allows patients and providers to make healthcare decisions together’. Treatment adherence, HIV testing, psychological care and sexual risk were the main areas where SDM was undertaken. These link closely to the main decision areas reported by patients in the review of patient decision making by Bravo *et al.* [1]. However, Bravo *et al.* also highlighted ‘decisions about disclosure to others’ and ‘decisions about starting a family’ as a key issues for patients [1]. SDM in these areas was reported much less frequently in our sample (Q.10) indicating that it may be useful to compare patient perspectives of SDM with that of professionals.

In spite of the apparent agreement in definition of SDM found in the quantitative data, the qualitative findings in this study highlight several nuances around how SDM is understood and practised. For many nurses in the group discussions, SDM was perceived as a broad issue of ‘involving the patient in care’, rather than as a very specific process around specific decisions as recommended in the literature/guidance in this area. Reflecting the holistic perspective of nursing, for many participants, SDM was about a decision ‘in context’ and a decision ‘within a relationship’ and much less about the mechanics and steps of the actual decision-making process, which were more vaguely described. Indeed, the most commonly described strategies for SDM were ‘general discussion’ rather than following any specific framework, for example, the Ottawa Decision Framework [22]. Hence, whilst the literature on SDM defines it as quite a narrow and specific phenomenon [6], nurses appeared to have a more inclusive but, arguably, hazy view of it. However, this may also reflect the fact that the majority of participants completing the questionnaire (86%) reported having received very little formal training on SDM (Q.14).

Another issue that emerged from the qualitative data was that nurses articulated a collaborative, inclusive and team-based understanding of SDM. Much of the discussion referred to nurses’ relationships with other health professionals and their role in representing the patient in discussions with other health professionals and within MDT meetings. In this way, nurses seemed to be acting indirectly as **facilitators** of SDM, as well as directly engaging in SDM with patients themselves. This facilitation role was also evident when nurses described themselves as ‘translating’ between doctors and patients to ensure that each understood the other’s perspective. The role that nurses play in facilitating SDM has not been explicitly recognised in the literature around SDM, which has tended to focus very much on the relationship and communication between a particular professional–patient dyad [4]. In contrast, our study suggests that shared models of care and MDT meetings represent key sites for decision making and that nurses play an important role in advocating within these for greater patient involvement.

Finally, the study has also revealed that SDM can be challenging for nurses. Some of these challenges are related to patients. Nurses experienced a tension in adopting an SDM approach with patients perceived to be complex, chaotic or who were making choices contrary to current guidelines. Similarly, the questionnaire data showed that nurses perceived some patients as not wanting to engage in SDM at all, preferring the nurses to advise them on what to do (Q.12 and Q.13). The most significant challenge of all was posed by patients who wished to opt out of treatment altogether. The questionnaire data showed that nurses would appreciate greater training around SDM, with a specific focus on how to support these more complex patient scenarios (Q.15). Other challenges around SDM were more structural. They related to the changes that were occurring within the disease area, which required participants to adhere to national targets, guidelines and protocols, all of which served to inhibit patient choice and therefore limit the opportunity to engage in SDM in a meaningful way. Lastly, there were organisational challenges. Nurses noted that SDM required more time, which was often in short supply, and more resources. With regard to the latter, the questionnaire indicated that 72% of the respondents wanted greater access to resources to facilitate SDM (Q.17) and 52% said that treatment decision aids would help them to implement

SDM (Q.9). The need for more resources is also reflected in Q.8 where only 12.5% said they 'always used' decision aids with patients.

Limitations of the research

The research explored SDM from the perspective of the HIV nurse and did not take into consideration the views and perspectives of patients, or those of other health professionals involved in HIV care. As such, the interpretations offered here are limited to one professional perspective.

Recommendations

The issues raised by this report can be addressed in three ways:

1. **Education/training for professionals.** There is a clear need for additional training to be provided for HIV nurses around SDM, with a specific focus on supporting challenging patient situations. The questionnaire (Q.17) indicated that 78% of respondents wanted greater knowledge about SDM and 56% wanted this training to be 'practical, including role playing'. The majority of respondents wanted training to be delivered as e-learning or study days.
2. **Development of decision support resources.** There is a need to develop decision support aids that nurses can use with their patients on a range of topics. This study has clearly identified a gap in availability of resources to support SDM.
3. **Further research.** In order to inform the content of training or resources, it is important to understand the patient perspective for SDM. There is currently a lack of research on patient experiences of SDM in UK HIV care. Such research is required. Similarly, undertaking research on the views of other health professionals around SDM would help to provide a more comprehensive picture of current issues and challenges.

Conclusions

This study has demonstrated HIV nurses' commitment to SDM. It has highlighted hitherto under-recognised elements of the nurse's role in SDM in terms of facilitating the process within a wider MDT. The study has also highlighted several patient-related, organisational and structural challenges to SDM. Finally, the study has identified a need for greater training for nurses around SDM and for the development of decision-support resources. As the treatment of HIV infection progresses towards a chronic disease-management model, developing skills in SDM is essential in order to facilitate better long-term health outcomes.

Acknowledgements

The team thanks Gilead Sciences Ltd for providing funding for this project. We are grateful to Natasher Lafond for helping with the research process. Finally, we thank all the participants for kindly giving up their time to share their views and experiences on shared decision making.

Abbreviations

ART	Antiretroviral therapy
ARV	Antiretroviral
FGD	Focus-group discussion
HIV	Human immunodeficiency virus
MDT	Multidisciplinary team
NHIVNA	National HIV Nurses Association
SDM	Shared decision making

References

1. Bravo P, Edwards A, Rollnick S, Elwyn G. Tough decisions faced by people living with HIV: a literature review of psychosocial problems. *AIDS Rev* 2010; **12**: 76–88.
2. Coulter A, Collins A. *Making shared decision making a reality. No decision about me without me*. London: The King's Fund; 2011. Available at: http://www.kingsfund.org.uk/sites/files/kf/Making-shared-decision-making-a-reality-paper-Angela-Coulter-Alf-Collins-July-2011_0.pdf (accessed January 2016).
3. Department of Health. *No decision about me without me: liberating the NHS. Government response*. London: HMSO; 2012. Available at: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/216980/Liberating-the-NHS-No-decision-about-me-without-me-Government-response.pdf (accessed January 2016).
4. Lee O, Emmanuel E. Shared decision making to improve care and reduce costs. *N Engl J Med* 2013; **368**: 6–8.
5. Friesen-Storms JH, Bours GJ, van der Weijden T, Beurskens AJ. Shared decision making in chronic care in the context of evidence based practice in nursing. *Int J Nurs Stud* 2014; **52**: 393–402.
6. Stiggelbout AM, Pieterse AH, De Haes JC. Shared decision making: concepts, evidence, and practice. *Patient Educ Couns* 2015; **98**: 1172–1179.
7. National AIDS Trust. *No decision about me without me: consultation on shared decision making. Response from National AIDS Trust, NAT*. London: National AIDS Trust; 2012. Available at: <http://www.nat.org.uk/media/Files/Policy/2012/Aug-2012-Shared-decision-making-NHS.pdf> (accessed January 2016).
8. British HIV Association. *Standards of Care for People Living with HIV 2013*. London: BHIVA; 2012. Available at: <http://www.bhiva.org/documents/Standards-of-care/BHIVASStandardsA4.pdf> (accessed January 2016).
9. European AIDS Clinical Society. *EACS Clinical Guidelines. Version 7.1*. Brussels: European AIDS Clinical Society; 2014. Available at: www.eacsociety.org/files/guidelines_english_71_141204.pdf (accessed February 2016).
10. Legare F, Ratte S, Gravel K, Graham ID. Barriers and facilitators to implementing shared decision-making in clinical practice: update of a systematic review of health professionals' perceptions. *Patient Educ Couns* 2008; **73**: 526–535.
11. Shay LA, Lafata JE. Understanding patient perceptions of shared decision making. *Patient Educ Couns* 2014; **96**: 295–301.
12. Truglio-Londrigan M, Slyer JT, Singleton JK, Worrall PS. A qualitative systematic review of internal and external influences on shared decision-making in all health care settings. *JBI Database of Systematic Reviews & Implementation Reports* 2014; **12**: 121–194.
13. Kumar R, Korthuis PT, Saha S *et al*. Decision-making role preferences among patients with HIV: associations with patient and provider characteristics and communication behaviors. *J Gen Intern Med* 2010; **25**: 517–523.
14. Laws MB, Beach MC, Lee Y *et al*. Provider–patient adherence dialogue in HIV care: results of a multisite study. *AIDS Behav* 2013; **17**: 148–159.
15. Beach MC, Roter DL, Saha S *et al*. Impact of a brief patient and provider intervention to improve the quality of communication about medication adherence among HIV patients. *Patient Educ Couns* 2015; **98**: 1078–1083.
16. Creswell J, Plano-Clark V. *Designing and Conducting Mixed Methods Research*. Thousand Oaks, California: Sage; 2007.
17. Barbour R. *Doing Focus Groups*. London: Sage Publications; 2007.
18. Richards L. *Handling Qualitative Data: A Practical Guide*. London: Sage; 2009.
19. Bryman A, Burgess R. *Analysing Qualitative Data*. London: Routledge; 1994.
20. Boyatzis R. *Transforming Qualitative data: Thematic Analysis and Code Development*. London: Sage; 1998.
21. Lincoln Y, Guba E. *Naturalistic Inquiry*. California: Sage; 1985.
22. Legare F, O'Connor AM, Graham ID *et al*. Primary health care professionals' views on barriers and facilitators to the implementation of the Ottawa Decision Support Framework in practice. *Patient Educ Couns* 2006; **63**: 380–390.

Appendix: online survey

Preamble

Dear NHIVNA member

Shared decision making is an emerging topic in HIV nursing care at the moment and NHIVNA wishes to gain insight into your views and experience of this issue. We would also like to establish if there is a need for further education and support on this topic.

In order to gather accurate and representative information we would be very grateful if you could complete this questionnaire, which should take up to 15 minutes to complete.

The findings will be analysed, published and used to develop resources to support your educational and clinical requirements. The questionnaires are anonymised unless you chose to give your name and contact details at the end.

Questions

Demographics

Please indicate the setting in which you currently work. (Please tick all that apply.)

- Community
- Specialist HIV Outpatients Department
- Genitourinary Medicine
- Midwifery/Obstetrics
- Infectious Diseases Unit
- Paediatrics
- Specialist HIV Inpatient Care
- Palliative/Hospice care
- Voluntary sector
- Other (please indicate)

Which of the following geographical areas do you work in? (Please tick one)

- England
- Ireland
- Scotland
- Wales

What is the approximate size of your HIV patient cohort in your work setting?

- Up to and including 500
- Over 501

Please indicate at which level you are currently employed?

- Band 5
- Band 6
- Band 7
- Band 8
- Other or non NHS, please provide details

How long in total have you been working specifically in HIV care?

- Not specialising in this field
- Less than 1 year

- 1–2 years
- 2–5 years
- 5–10 years
- More than 10 years

Shared decision making

In your view, which of the following statements **best defines shared decision making** in HIV nursing care? (Please tick one.)

- Informing the patient of the best treatment options available
- A process to help engage with a patient when they have disengaged or stopped treatment
- Checking what the patient understands and how the patient feels about what you've said
- Being an advocate for patients when you know what they want
- Planned care to which the patient consents
- Shared care management with other health professionals
- Giving the patient all the necessary information so that they can make the decision on their own
- Multidisciplinary teams working together to make the best healthcare decisions for the patient
- A collaborative process that allows patients and their providers to make healthcare decisions together
- Shared decision making is not possible because patients are always asking me what I would do

On a scale of 1 to 5, please indicate **how confident you feel** in undertaking shared decision making in the following areas of care where 1 is not confident at all and 5 is very confident.

- HIV testing
- Starting HIV treatment
- Switching HIV treatment
- Stopping HIV treatment
- Opting out of treatment
- Adhering to treatment
- Smoking cessation
- Sexual risk taking
- Family/pregnancy planning
- Recreational drug use
- Managing HIV treatment along with treatment for other medical conditions
- Psychological care
- Disclosure of status to others
- Dietary advice
- Exercise advice
- Cardiovascular risk advice
- Alcohol consumption advice

On a scale of 1 to 5 how easy is it to practise shared decision making in your area of HIV care?

- Very easy
- Easy
- Moderate
- Hard
- Very Hard

In your day-to-day practice, in which activities do you **most commonly undertake** shared decision making with your client? Please give your answer on the scale of 1 to 5 where 1 indicates 'very common' and 5 indicates 'very rarely'.

- HIV testing
- Starting HIV treatment
- Switching HIV treatment
- Stopping HIV treatment
- Opting out of treatment
- Adhering to treatment
- Smoking cessation
- Sexual risk taking
- Family/pregnancy planning
- Recreational drug use
- Managing HIV treatment along with treatment for other medical conditions
- Psychological care
- Disclosure of status to others
- Dietary advice
- Exercise advice
- Cardiovascular risk advice
- Alcohol consumption advice

Have you had any specific training on the topic of 'shared decision making'? Please state:

- Yes (if yes, please state)
- No
- Unsure

What strategies do you most use to facilitate shared decision making? Please rank on a scale of 1 to 5 where 1 = never use, 2 = not very often but have used, 3 = occasionally use, 4 = frequently use, 5 = always use.

- General discussion
- Motivational interviewing
- Use of written decision aids
- Use of web-based resources
- Drawing on your own professional knowledge/experience
- Advocacy with other services or professionals
- Signposting to other services
- Linking to peer support
- Other; please list

The following have been identified as potential barriers to implementing shared decision making in everyday consultations. Please tick the top three that have applied to you in your clinical role.

- Not enough consultation time
- Feeling unskilled
- Lack of up to date background knowledge on the patient
- Lack confidence in my own skills
- Lack of support from colleagues
- This is the role of another member of the MDT
- Needing interpreters
- Cultural issues/patients beliefs
- The setting doesn't support shared decision making
- Treatment guidelines/options available
- Not knowing patient adequately
- Organisational targets
- Public Health England targets
- Patient doesn't want to input into the process/patient want me to make the decision
- Other barriers; please describe

What would help you to implement shared decision making in your day-to-day care? Please tick your top three.

- Better background knowledge
- Treatment decision aids
- More training in communication skills
- More educational resources for patients
- More time with patients in consultations
- More support from other professionals
- More support from the voluntary sector
- None of the above
- Other; please describe

Please select the top three issues from the options below where greater knowledge/skills would help to enhance your ability to practise shared decision making in HIV care.

- Resources that address language barrier
- Patients who refuse treatment
- Patient non attenders
- Patients who take recreational drugs
- Patients with drug/alcohol addictions
- Patients with mental health problems
- Chaotic patients
- Assessing patients capacity to make treatment decisions

Which resources have you accessed to date, in order to enhance your knowledge of shared decision making? Please tick any that apply.

- Study days
- Conference lectures

- Online resources
- In-house education
- Medical/nursing journal articles
- Information booklets
- None
- Other; please describe

In your view, what knowledge/skills would enable you to undertake SDM more effectively with your clients? Please select the top three.

- Greater knowledge of SDM process
- Motivational interviewing
- Counselling skills
- Practical training in SDM process including role playing
- Access to resources to aid SDM

Are there any limitations of shared decision making? Please select all that apply.

- Not everyone wants shared decision making
- Shared decision making is not compatible with clinical practice guidelines
- Shared decision making takes too much time
- Not everyone is good at shared decision making
- There are no limitations to shared decision making
- Other; please describe

If further training and education were made available on the subject of shared decision making, how would you prefer to access this? Please give your order of preference, where 1 would be your first choice.

- e-learning
- Reading materials online
- Hard copy reading materials
- Study days
- Conference lectures
- Interactive workshops at conferences
- Short education sessions run in my workplace
- Other suggestions; please describe. ...

Now we are almost finished

The survey is completely anonymous (but there is an option to provide us with an email address so that we could enter you into a prize draw for a chance to win £100 worth of Amazon vouchers). If you would like to be entered into the Amazon voucher prize draw, please provide your email address here:

.....

Many thanks for taking the time to complete this survey. The results and outcomes will be disseminated through NHIVNA activities during 2015.

2016



Exploring shared decision making in HIV nursing care

NHIVNA aims to provide an academic and educational forum for the dissemination of original nursing research in the field of HIV/AIDS. We aim to address the communication and support needs of nurses working in this area.

We hope that these activities will assist in the promotion of good practice in the care of people with HIV.

Published by: **Mediscript Ltd**
© NHIVNA 2016

NHIVNA Secretariat

1 Mountview Court · 310 Friern Barnet Lane · London N20 0LD
Tel: +44 (0)20 8446 8898 · Fax: +44 (0)20 8446 9194 · Email: nhivna@mediscript.ltd.uk · Web: www.nhivna.org

Registered Charity No: 1099074